

Managing Phytophthora Root and Stem Rot of Soybeans

by Steve Butzen and Debra Steiger

Summary

- Phytophthora is a disease of extended wet field conditions. Such conditions may occur in any field but are most common on heavy soils.
- Phytophthora may attack soybeans at any time during the growing season. The disease displays a seed rot phase, a seedling blight phase, and a root and stem rot phase.
- Selecting varieties with specific race resistance or field tolerance is the primary means of combating Phytophthora.
- Varieties with specific race resistance provide a high level of protection, but may lose effectiveness if Phytophthora races change. Tolerant varieties provide protection against all races of Phytophthora.
- Other management practices include improving field drainage, remediating compaction and hardpans, planting fields when soils are warm, and using seed treatments.

Phytophthora is a soybean disease associated with wet soil conditions commonly occurring on heavy, poorly-drained or compacted soils. However, if normally well-drained fields are saturated for an extended period due to heavy rainfall or even excessive irrigation, Phytophthora can infect those fields as well. The ideal temperature for infection is 60 to 80F. Disease severity is highly dependent on environmental conditions, Phytophthora races in the field, and genetic resistance or field tolerance of the soybean variety. Depending on those factors, yield reductions can range from as little as 5% to over 50% when infection occurs in a field. When early infection results in severe seed rot and damping off of seedlings, replanting is a common outcome. This *Crop Insights* will describe the stages and symptoms of Phytophthora on soybeans, and how to minimize its effects by use of genetically resistant and field tolerant varieties, and employing other management practices.

Phytophthora Development and Symptoms

Phytophthora may attack soybeans at any time during the growing season. The seed rot phase may affect soybeans as soon as seeds imbibe water and swell. The seedling blight phase may occur prior to emergence or during early seedling growth. The root and stem rot phase may occur any time throughout the summer.

Seed Rot Phase – Soybeans seeds are at risk to Phytophthora as soon as seeds begin the germination process. Infected seeds are dark brown and soft to mushy. Severe infection at this stage may result in complete deterioration of the seed. Seed rot is most common when soils become saturated immediately after planting, conditions remain wet, and temperatures are optimum for disease development.

Seedling Blight Phase - The seedling blight phase of Phytophthora may infect plants during the emergence process or shortly thereafter. Severe infections can result in rapid decay, wilting and death of the just-emerged seedling, commonly referred to as “damping off”. Symptoms include a dark brown to black discoloration of the stem, usually beginning at the soil line. Diseased tissues quickly become soft and water-soaked, and wilting and plant death may soon follow. Seedling plants may also succumb to Phytophthora at later stages.

Another soil pathogen, Pythium, can also result in seed rot and damping off. Pythium is the more active of these two pathogens from 50 to 60F. Above 60F, Phytophthora is more active.



Dark-brown lesion on lower stem is a key symptom of Phytophthora¹.

Root and Stem Rot Phase

- Because the Phytophthora fungus is a soil-borne organism, infections begin in the root of the plant and then progress into the lower stem. Infections may eventually kill the soybean plant outright or may just weaken it to varying degrees. Even if plants survive, productivity is often greatly reduced compared to that of healthy soybean plants.

Root symptoms include brown, discolored secondary roots and an overall reduction in root mass that allows the plant to be easily pulled from the ground. The main tap root is also brown on the surface, and splitting the root reveals a tan discoloration of the inner tissues. Nodulation is often minimal on the roots and this is expressed as light-green plants above ground. Plants may also be stunted,

resulting in fields with an uneven appearance. Root damage results in wilting of the plant during periods of stress.

If the root infection spreads into the stem, a brown discoloration begins to develop at the soil line. Eventually, a dark-brown to red-brown lesion may progress several nodes up the stem from the base of the plant (see image on page 1). This girdling lesion is one of the key diagnostic features of the stem rot phase of Phytophthora. However, disease progression may not reach this stage when infection is moderate or begins later in the season.

Above-ground symptoms of Phytophthora may not be evident for several weeks after initial infection. Often, a period of drought stress is required before diseased plants begin to show the effects of reduced root systems and inadequate water transport capability within the stem. Symptoms progressing from the bottom to the top of the plant include leaves wilting followed by petioles drooping. As plants reach the permanent wilt stage, leaves may develop a grayish cast followed by yellow spots.

Management Practices for Phytophthora

Variety Selection – Phytophthora is a highly diverse organism – over 70 different races have been identified throughout the US. In a given field, many different races are represented in the Phytophthora population, with one or two of these being the most dominant (most frequently occurring). A field population of Phytophthora is dynamic – it is constantly changing in response to climate and soil conditions, varieties grown, crop rotation and other factors.

Genetic resistance bred into soybean varieties is race-specific. This means it is able to defeat certain races, but is overcome by others. When the dominant race in a field shifts, it may overcome the resistant soybean varieties developed for an area. Soybean breeders must then find a new source of resistance to defeat that race, and incorporate it into new varieties. This illustrates the drawback of race-specific resistance – it may eventually lose its effectiveness as Phytophthora populations change. In fact, use of the same resistance gene continually will eventually cause a race shift that reduces the effectiveness of the gene.

Another type of genetic protection against Phytophthora is known as “field tolerance”. This term is alternately referred to as “tolerance”, “partial resistance”, and “rate-reducing resistance” by various researchers and seed companies. Field tolerance is not race-specific – it is effective against all races of Phytophthora. This type of protection is more enduring over time than that offered by specific resistance genes, which may eventually need to be replaced.

Varieties containing both genetic resistance and field tolerance have two mechanisms of protection. In areas where a resistance gene is beginning to lose its effectiveness, the additional protection afforded by the field tolerance trait is very useful. A difference in field tolerance is often the reason

that two competing varieties, each with the same resistance gene, may perform very differently in the field.

The effectiveness of several common resistance genes to 25 races of Phytophthora is shown in Table 1. (Phytophthora races differ from area to area, and new races emerge as the most dominant in an area. For that reason, the races indicated here may not be the most important ones for your area.) Pioneer researchers evaluate soybean varieties in hundreds of environments throughout North America. These researchers are developing new varieties with the appropriate resistance genes for each area, as well as incorporating higher levels of field tolerance into new varieties. Your local Pioneer sales professional can help you select the appropriate Phytophthora resistant or tolerant varieties for your fields.

Table 1. Effectiveness of several Phytophthora resistance genes against various races of Phytophthora (green = effective.)

Race of Phytoph.	----- Resistance Gene (Rps) -----						
	1a	1b	1c	1k	3	4	6
1	Green	Green	Green	Green	Green	Green	Green
2	Green	White	Green	Green	Green	Green	Green
3	Green	Green	Green	Green	Green	Green	Green
4	Green	Green	Green	Green	Green	Green	Green
5	Green	Green	Green	Green	Green	Green	Green
6	Green	Green	Green	Green	Green	Green	Green
7	Green	Green	Green	Green	Green	Green	Green
8	Green	Green	Green	Green	Green	Green	Green
9	Green	Green	Green	Green	Green	Green	Green
10	Green	Green	Green	Green	Green	Green	Green
11	Green	Green	Green	Green	Green	Green	Green
12	Green	Green	Green	Green	Green	Green	Green
13	Green	Green	Green	Green	Green	Green	Green
14	Green	Green	Green	Green	Green	Green	Green
15	Green	Green	Green	Green	Green	Green	Green
16	Green	Green	Green	Green	Green	Green	Green
17	Green	Green	Green	Green	Green	Green	Green
18	Green	Green	Green	Green	Green	Green	Green
19	Green	Green	Green	Green	Green	Green	Green
20	Green	Green	Green	Green	Green	Green	Green
21	Green	Green	Green	Green	Green	Green	Green
22	Green	Green	Green	Green	Green	Green	Green
23	Green	Green	Green	Green	Green	Green	Green
24	Green	Green	Green	Green	Green	Green	Green
25	Green	Green	Green	Green	Green	Green	Green

Source: South Dakota State University, 2001. (Note that races shown here may not be the most important ones for your area.)

Field Drainage - Because Phytophthora development requires saturated soils for a prolonged period, improving field drainage is one key to successful management. Other field characteristics that contribute to waterlogged soils include compaction and hardpan layers. Remediating these soil

structure issues is another useful management tool to reduce Phytophthora infections.

Planting Date – Early planting may increase the risk of Phytophthora damage if soils are wetter and cool temperatures slow soybean emergence. Planting later when temperatures are more favorable to soybean growth gives soybeans the edge in outpacing disease development. This is more important on heavy soils or in no-till farming systems where seedbeds are intrinsically cooler.



Soybean seedlings dying under high Phytophthora pressure. Seed treatments help provide early protection.

head start on the diseases. Fungicide seed treatments are especially useful when cool, wet soil conditions develop after planting. These conditions favor diseases and slow soybean growth, creating an ideal scenario for crop damage. Metalaxyl and mefenoxam are available in a number of formulations of Apron® and Allegiance® fungicide seed treatments. Read and follow label instructions carefully, including rates required for Phytophthora control.

Manure and Fertilizer Application – Application of high levels of potash, manure or municipal sludge immediately before planting soybeans may result in more severe Phytophthora root rot (Dorrance, et al., 2002) This is because of the chloride, salt or nitrate contained in fertilizer, manure and sludge. Application of these products should be made in the fall to allow time for leaching of soluble salts prior to planting. Growers should also avoid concentrating manure or sludge in specific fields, even if they are closer to the source.

Seed Treatments – In some fields with a history of Phytophthora damage, seed-applied fungicides should be routinely considered. Two fungicide active ingredients – metalaxyl and mefenoxam – have specific activity against Phytophthora and Pythium. These fungicides may provide protection for up to two to three weeks. This early period of protection often covers the period from planting to emergence and allows soybean seedlings a

Conclusion

Resistant and field tolerant varieties provide the first line of defense against Phytophthora. Pioneer Hi-Bred has resistant varieties in most maturity groups, as well as varieties with excellent field tolerance to this disease. In addition, the highest quality seed with high germination and good seedling vigor can improve stand establishment and overall performance in the presence of Phytophthora.

References

- Dorrance, A., P. Lipps and D. Mills. 2002. Phytophthora damping off and root rot of soybean. Extension Factsheet A-17-02. The Ohio State University.
<http://ohioline.osu.edu/ac-fact/0017.html>
- Draper, M and T. Chase. 2001. Phytophthora root and stem rot (PRR) of soybean. Extension Fact Sheet 902-B, South Dakota State University.
<http://agbiopubs.sdstate.edu/articles/FS902B.pdf>
- Hershman, D. and P. Bachi. 2003. Phytophthora root and stem rot of soybean. Plant Pathology Factsheet PPFs-AG-S-4. University of Kentucky.
http://www.ca.uky.edu/agcollege/plantpathology/ext_files/PPF_Shtml/ppfsags4.pdf

¹ Image courtesy of Jim Boersma, Pioneer Area Agronomist.

² Image courtesy of Viarural: www.viarural.com.ar.

Apron is a registered trademark of a Syngenta Group Company.

Allegiance is a registered trademark of Bayer AG.